

## Seasonal variation in suicides: diminished or vanished

Experience from England and Wales, 1982–1996

PAUL S. F. YIP, ANNE CHAO and CALVIN W. F. CHIU

**Background** Seasonal variation in suicidal death has been observed in many countries. In particular, a cyclic variation was found for both men and women in England and Wales in the 1960s and 1970s. Men showed a single 12-month cycle whereas women showed two cycles.

**Aims** To re-examine the seasonal variation in suicides in England and Wales for the period 1982–1996.

**Method** A harmonic analysis was used to detect the seasonality of the suicide data.

**Results** The seasonal effect on suicide is greatly diminished in England and Wales. This is shown by the reduced amplitude and smaller proportion of variance accounted for by the season.

**Conclusions** The seasonal effect on suicide has either diminished or vanished.

**Declaration of interest** None.

Seasonal variation in deaths by suicide is an important study in terms of understanding the possible sociological and biological determinants of suicide and its prevention. Spring peaks for males and spring and autumn peaks for females were found in the UK (Barraclough & White, 1978; Meares *et al*, 1981), Australia (Eastwood & Peacocke, 1976; Parker & Walter, 1982), Finland (Nayha, 1982, 1983) and Italy (Micciolo *et al*, 1989, 1991). A spring and late summer peak and a trough in winter months were found for suicides for both men and women in the USA (Lester, 1971; Lester & Frank, 1988). Massing & Angermeyer (1985) give a comprehensive account of the effects of season on suicides in the 1960s and 1970s. Recently, a harmonic analysis was applied to examine suicides in Australia and New Zealand. Only one cycle for both men and women was observed (Yip *et al*, 1998). A similar finding was also observed in Hong Kong and Taiwan (Ho *et al*, 1997). The present study attempts to examine seasonal effects on suicides in England and Wales with respect to gender, age group and method of suicide. We have two objectives: to verify the findings in Meares *et al* (1981) and Barraclough and White (1978) about the seasonal effect in England and Wales, i.e. men showed a single 12-month cycle, whereas women showed two cycles; and to investigate whether seasonal variation is found more strongly in some subgroups of the population (e.g. by age or method of suicide) than in others.

### METHOD

The Office for National Statistics (England & Wales) provided data for all deaths by suicide for the period 1982–1996. Information on age, gender, method and month of suicide was also made available. The data included here relate to cases where judicial inquiries established that the cause

of death was suicide. They were coded in the range E950–959 of ICD-10 (World Health Organization, 1992). The chosen period enabled comparison with results obtained in the 1960s and 1970s (Barraclough & White, 1978; Meares *et al*, 1981).


The seasonal variations in suicides were examined in three ways. First, the number of suicides in England and Wales for each month of the period 1982–1996 was plotted separately for each gender. Second, a daily mean suicide incidence and cumulative number of suicides were calculated for each month of the study period. Third, a harmonic time series model was applied to different genders, age groups and methods of suicide. The method of analysis was similar to that employed by Barraclough & White (1978), Micciolo *et al* (1991), Ho *et al* (1997) and Yip *et al* (1998). In this model, the variation in suicides between the months is described as the sum of the sinusoidal curves. The seasonal variation consists of those components with cycles that repeat themselves an exact number of times per year. Let  $A_i$  be the number of suicides for month  $i$ ; for a period of 15 years (180 months),

$$A_i = a_0 + \sum_{j=1}^{90} \left( a_j \cos \frac{2\pi ij}{180} + b_j \sin \frac{2\pi ij}{180} \right) \\ i=1, \dots, 180$$


where  $a_j$  and  $b_j$  are constant ( $j=1, \dots, 90$ ),  $a_0$  is the mean suicide number and

$$a_j \cos \frac{2\pi ij}{180} + b_j \sin \frac{2\pi ij}{180}$$

is commonly referred to as the  $j$ th harmonic of  $A_i$  with period  $15/j$  years and frequency per annum  $j/15$ . The harmonics with  $j=15, 30, 45, 60, 75$  and  $90$  have periods of 1, 1/2, 1/3, 1/4, 1/5 and 1/6 years and frequencies per annum of 1, 2, 3, 4, 5 and 6, respectively. For example, one such sinusoidal curve is that with a period of 6 months (1/2 year), which has just two peaks and two troughs in each year. These harmonics have a cycle that repeats an exact number of times per year and follows the same pattern in years as any combinations of such harmonics. The quantities  $a_j$  and  $b_j$  are estimated so as to give the best fit to the data, and they describe the amplitudes of the separate sinusoidal components. The significance of a particular



**Fig. 1** Age-specific suicide rates (per 100 000) in England and Wales, 1982–1996: —●—, male 60+; —▲—, male 40–59; —×—, male 25–39; —+—, male 15–24; - -●- -, female 60+; - -▲- -, female 40–59; - -×- -, female 25–39; - -+ - -, female 15–24.



**Fig. 2** Monthly distribution of suicide deaths in England and Wales, 1982–1996.

component  $j$  is ascertained by testing whether  $a_j$  and  $b_j$  are significantly different from zero; if they are not, it may be concluded that there is no seasonal variation. It is assumed that the total variance of the

distribution of the monthly suicide data can be broken down into three components: random, seasonal and non-seasonal. In this way it is possible to calculate the percentage of total variance attributable to

seasonal variation, as well as to random and non-seasonal variation. Under the alternative hypothesis, that the variation is purely random, the monthly totals may be considered as independent identically distributed Poisson random variables. The details of testing the significance of the different components of the variation can be found in Pocock (1974), Barraclough & White (1978) and Yip *et al* (1998).

**RESULTS**


A total of 59 608 (43 229 males and 16 379 females) suicides were identified in England and Wales for the period 1982–1996. The age-specific suicide rates by gender for the study period were plotted and are shown in Fig. 1. A statistically significant decreasing trend was observed in the group (male and female) aged 60 years or over ( $P < 0.05$ ). On the other hand, an increase in the suicide rate among males of age 15–24 years was noted ( $P < 0.05$ ). Suicide rates increased with age among females, but not among males. Males aged 25–39 years have shown the highest rate since 1994. The monthly distribution of suicides, for males and females is shown in Fig. 2. The cumulative number of suicides for both males and females, and the monthly mean number of suicides per day are shown in Table 1 and Fig. 3, respectively. The assumption of even distribution of suicides was rejected ( $P < 0.01$ ): December had a trough for both males and females.

Table 2 shows the results of harmonic analysis by gender. Only 15 and 17% of the total variations can be explained by the seasonal component for males and females, respectively. The effects of all seasonal harmonics are marginally statistically significant, with  $P \approx 0.05$ . In order to examine the possible determinant

**Table 1** The observed and expected number of suicides by month and gender in England and Wales, 1982–1996

	January	February	March	April	May	June	July	August	September	October	November	December
<b>Male</b>												
Observed	3755	3251	3777	3706	3717	3660	3669	3626	3481	3590	3605	3392
Expected <sup>I</sup>	3669	3345	3669	3550	3669	3550	3669	3669	3550	3669	3550	3669
$\chi^2=43.98, d.f.=11, P < 0.0001$												
<b>Female</b>												
Observed	1362	1244	1496	1452	1448	1376	1370	1301	1337	1351	1416	1226
Expected <sup>I</sup>	1390	1268	1390	1345	1390	1345	1390	1390	1345	1390	1345	1390
$\chi^2=50.90, d.f.=11, P < 0.0001$												

I. The expected number of deaths has been adjusted by the number of days in a month.



**Fig. 3** Monthly averages of suicide incidence over 15 years in England and Wales, 1982–1996.

factors of season by different subgroups, the figures for age and methods of suicide were also subjected to the same analysis. The results obtained on the seasonal components (not shown here) are not statistically significant.

## DISCUSSION

The present study suggests that the marked seasonal fluctuation in suicide rates found in England and Wales in the 1960s and 1970s appears to be diminishing, and in some cases it has vanished completely (Barraclough & White, 1978; Meares *et al*, 1981). Barraclough & White (1978) used the same method as described here. They examined the seasons during the period between 1968 and 1972 and found that 49.3% of the variation could be explained by season, with the adjusted ratio 2.5:1 of the components of all seasonal harmonics to random. The effect of season on suicide during the period 1982–1996 accounts for only 15–17% of the variation. The results suggest, therefore, a

significant reduction in the proportion of variance accounted for by season. We examined the possible confounding factors, such as age and method of suicide, and found nothing to be significant. Also, we included the undetermined causes of death (E980–989) in the analysis and similar results were obtained. Seasonal components were not significant.

### Reduction in seasonal fluctuation observed in other countries

Similar findings using the same harmonic analysis were obtained in other countries: for example, Micciolo *et al* (1989) suggested that 48–65% of the variance in Italy's suicide data in the 1970s was explained by seasonal components; in the 1980s and 1990s this figure was only 25–32% in Hong Kong and Taiwan (Ho *et al*, 1997), and 3–17% in Australia and New Zealand (Yip *et al*, 1998).

### Significant reduction in suicides among the widowed and divorced

The reduction of the seasonal component can be explained by the significant reduction in suicides among the widowed and divorced in England and Wales for the past two decades. In fact, the decrease in suicide rates in England and Wales from the widowed and divorced groups accounts for most of the overall fall in the suicide rate for men and women. For example, the suicide rates for males and females aged 60 years or over have fallen from 18.2 to 11.1 and from 10.9 to 4.0, respectively, for the period 1982–1996. The precise reasons for the decrease among the

widowed and divorced, especially among the elderly, are yet to be determined. It has been shown that in Finland the autumn peak is related to marital status. There was a high frequency of suicides among the divorced (and possibly widowed) in September. Furthermore, divorced males, who are known to be located in a lower than average position in the social scale, might be claimed to be liable to suicide in autumn, owing to the increased unemployment rate at that time of the year (Nayha, 1983; Yip, 1997). Reduction of suicide for the two groups would have an impact on the seasonal variations in suicides.

### Change of lifestyle

Durkheim (1897) believed that seasonal variation in suicidal behaviour was determined by the intensity of communal life and activities. Technological development in telecommunications and the fact that people tend to be 'connected' more often than before (for example, by mobile phones, e-mail and the internet) and also the different activities and social contacts nowadays, all may play a part in determining the seasonal effect on suicide.

The exact impact of these changes on the seasonal variation in suicides is far from clear, but it is not surprising to see a reduction in the seasonal influence on the distribution of suicides.

In conclusion, the present study found that the seasonal effect on suicides in England and Wales is less obvious than previous studies suggested. The present finding challenges the existence of seasonal effects and predicts that seasonal variation in suicides will disappear in the new millennium.

**Table 2** Harmonic analysis of monthly data on suicides in England and Wales, 1982–1996

Component of variance	Male		Female	
	Value	Percentage	Value	Percentage
All seasonal harmonics	49.54	15%	24.02	17%
1 cycle <sup>1</sup>	18.36	37%	7.21	30%
2 cycles	0.21	0%	6.55	27%
3 cycles	0.00	0%	0.88	4%
4 cycles	1.41	3%	0.00	0%
5 cycles	20.31	41%	2.95	12%
6 cycles	9.26	19%	6.42	27%
Non-seasonal harmonics	45.95	14%	23.58	17%
Random variations	240.16	72%	91.02	66%
Total variance	335.65	100%	138.62	100%

1. The percentage values represent the proportion of the variance of the all seasonal harmonics that can be explained by the respective cycles.

## ACKNOWLEDGEMENTS

We are grateful to James Thorburn and the three referees for the very useful suggestions and to the Office for National Statistics and the Office of Population Censuses and Surveys (England & Wales) for supplying the data for the present study. This work is supported by a CRCG grant from the University of Hong Kong.

## REFERENCES

Barraclough, B. M. & White, S. J. (1978) Monthly variation of suicide and undetermined death compared. *British Journal of Psychiatry*, **132**, 275–278.

**Durkheim, E. (1897)** *Le Suicide*. Paris: F. Alcan. English translation (1962) *Suicide. A Study in Sociology*. London: Routledge and Kegan.

**Eastwood, M. R. & Peacocke, J. (1976)** Seasonal patterns of suicide, depression and electroconvulsive therapy. *British Journal of Psychiatry*, **129**, 472–475.

**Ho, T. P., Chao, A. & Yip, P. S. F. (1997)** Seasonal variation in suicides re-examined: no sex difference in Hong Kong and Taiwan. *Acta Psychiatrica Scandinavica*, **95**, 25–31.

**Lester, D. (1971)** Seasonal variation in suicidal deaths. *British Journal of Psychiatry*, **118**, 627–628.

— & **Frank, M. L. (1988)** Sex differences in the seasonal distribution of suicides. *British Journal of Psychiatry*, **153**, 115–117.

**Massing, W. & Angermeyer, M. (1985)** The monthly and weekly distribution of suicide. *Social Science and Medicine*, **21**, 433–441.

**Meares, R., Mendelsohn, F. A. O. & Milgrom-Friedman, J. (1981)** A sex difference in the seasonal variation of suicide rate: a single cycle for men, two cycles for women. *British Journal of Psychiatry*, **138**, 321–325.

**Micciole, R., Zimmermann-Tansella, Ch., Williams, P., et al (1989)** Seasonal variation in suicide: is there a sex difference? *Psychological Medicine*, **19**, 199–203.

—, **Williams, P., Zimmermann-Tansella, Ch., et al (1991)** Geographical and urban–rural variation in the seasonality of suicide: some further evidence. *Journal of Affective Disorder*, **21**, 39–43.

**Nayha, S. (1982)** Autumn incidence of suicide re-examined: data from Finland by sex, age and occupation. *British Journal of Psychiatry*, **141**, 512–517.

— (1983) The biseasonal incidence of some suicides. Experience from Finland by marital status, 1961–1976. *Acta Psychiatrica Scandinavica*, **67**, 32–42.

**Parker, G. & Walter, S. (1982)** Seasonal variation in depressive and suicidal deaths in New South Wales. *British Journal of Psychiatry*, **140**, 626–632.

**Pocock, S. J. (1974)** Harmonic analysis applied to seasonal variations in sickness absence. *Applied Statistics*, **23**, 103–120.

**World Health Organization (1992)** *The Tenth Revision of the International Classification of Disease and Related Health Problems (ICD–10)*. Geneva: WHO.

## CLINICAL IMPLICATIONS

- Suicide risk was more uniformly distributed throughout the year than before, which might have implications for planning support services for those at risk of or attempting suicide.
- Sociocultural factors may play a more significant role in determining the seasonal effect in suicides.
- The decrease in suicides among the high-risk groups of the widowed and divorced is encouraging. It is important to understand the reason(s) in order to be able to reduce the suicide rates further in England and Wales.

## LIMITATIONS

- The results would be more statistically powerful if a longer series of data were available, although 15 years of data are quite sufficient.
- The effect of underreporting of suicides on the seasonal effect is yet to be determined.
- We were unable to determine the biological effect (e.g. seasonality in suicide mirroring seasonality in onset of biological depression and mania) on the reduction of seasonal variation of suicide.

PAUL YIP, PhD, Department of Statistics and Actuarial Science, The University of Hong Kong; ANNE CHAO, PhD, Institute of Statistics, National Tsing Hua University, Hsinchu, Taiwan; CALVIN CHIU, MSc, Department of Statistics, Harvard University, MA, USA

Correspondence: Paul Yip, Department of Statistics and Actuarial Science, The University of Hong Kong, Pokfulam Road, Hong Kong

(First received 6 September 1999, final revision 17 May 2000, accepted 18 May 2000)

**Yip, P. S. F. (1997)** Suicides in Hong Kong 1981–1994. *Social Psychiatry and Psychiatric Epidemiology*, **32**, 243–250.

—, **Chao, A. & Ho, T. P. (1998)** A re-examination of seasonal variation in suicides in Australia and Taiwan. *Journal of Affective Disorders*, **47**, 141–150.

# BJPpsych

The British Journal of Psychiatry

## Seasonal variation in suicides: diminished or vanished: Experience from England and Wales, 1982-1996

PAUL YIP, ANNE CHAO and CALVIN CHIU

*BJP* 2000, 177:366-369.

Access the most recent version at DOI: [10.1192/bjp.177.4.366](https://doi.org/10.1192/bjp.177.4.366)

---

### References

This article cites 13 articles, 6 of which you can access for free at:  
<http://bjp.rcpsych.org/content/177/4/366#BIBL>

### Reprints/ permissions

To obtain reprints or permission to reproduce material from this paper, please write to [permissions@rcpsych.ac.uk](mailto:permissions@rcpsych.ac.uk)

### You can respond to this article at

[/letters/submit/bjprcpsych;177/4/366](http://bjp.rcpsych.org/letters/submit/bjprcpsych;177/4/366)

### Downloaded from

<http://bjp.rcpsych.org/> on June 21, 2016  
Published by The Royal College of Psychiatrists

---